

My asthma triggers

Tick the triggers that make your asthma worse:

- | | |
|-----------------------------------|--|
| <input type="checkbox"/> Pollen | <input type="checkbox"/> Air pollution |
| <input type="checkbox"/> Exercise | <input type="checkbox"/> Other |
| <input type="checkbox"/> Cold/flu | (please list here): |
| <input type="checkbox"/> Stress | |
| <input type="checkbox"/> Weather | |

“ Always keep your blue reliever inhaler and your spacer with you. You might need them if your asthma gets worse. ”

Dr Andy Whittamore
Asthma + Lung UK's GP

Book your child's asthma review

You should book an asthma review at least once a year, or more if your child needs it. If your child has been to A&E, or been prescribed steroid tablets or liquid, you should book an asthma review straight away. Remember to bring:

- their action plan, to see if it needs updating
- any inhalers and spacers they have, to check they're using them in the best way
- their peak flow meter if they use one
- any questions about their asthma and how to manage it.

Date of next asthma review:

Healthcare professional contact details:

Asthma and Lung UK is a charitable company limited by guarantee with company registration number 01863614, with registered charity number 326730 in England and Wales, SC038415 in Scotland, and 1177 in the Isle of Man.



Parents and carers

As well as wheezing, coughing and a tight chest, your child might have their own unique symptoms that tell you their asthma is getting worse. You can list them here:

Does your child tell you when they need their asthma inhaler?

- Yes No

Does your child need help taking their asthma medicines?

- Yes No

Get the most from your child's action plan:

- take a photo and keep it on your phone, and on your child's phone if they have one
- stick a copy on your fridge door
- share your child's action plan with their school, holiday club or grandparents – whoever is caring for your child.

Learn more about what to do during an asthma attack:

www.asthmaandlung.org.uk/asthma-attacks

Get advice, support and information at **AsthmaAndLung.org.uk** or find us on social media:



Questions about asthma?

Talk to our friendly respiratory nurse specialists for more support.
Call **0300 222 5800**
(Monday to Friday, 9am to 1pm and 2pm to 5pm)

Last reviewed July 2025; next review July 2028. V1.



Child asthma action plan

Fill this in with your healthcare professional

This asthma action plan is for children who use separate preventer and reliever inhalers. If you are on a MART or AIR regime, please use our MART or AIR asthma action plan.

Name and date:

1 My daily asthma routine

I need to take my preventer inhaler every day.

My preventer inhaler is called:

and its colour is:

I take _____ puff(s)
in the morning.

I take _____ puff(s)
at night.

I rinse my mouth out afterwards. I do this every day even if my asthma's okay.

I must remember to use my spacer with my inhaler. If I do not have one, I'll go back to my healthcare professional and ask for one.

Other asthma medicines I take every day:

My reliever inhaler helps when I have symptoms.

My reliever inhaler is called:

and its colour is:

IMPORTANT: If I need my blue reliever inhaler when I do sports or activity, I need to see a healthcare professional.

2 When I feel worse

My asthma is getting worse if I have any of these symptoms:

- I wheeze, cough, my chest hurts, or it's hard to breathe
- I need my blue reliever inhaler 3 or more times a week
- I'm waking up at night because of my asthma (this is an important sign and I will book a next day appointment with my healthcare professional).

If my asthma gets worse, I will:

- take my preventer medicines as normal
- take _____ puff(s) of my blue reliever inhaler every 4 hours if needed
- see my healthcare professional within 24 hours if I do not feel better.

URGENT! If your reliever inhaler is not lasting 4 hours, you need to take emergency action now. **See section 3.**

Other things my healthcare professional says I need to do if my asthma is getting worse:

3 In an asthma attack

I'm having an asthma attack if I have any of these symptoms:

- my reliever inhaler is not helping or it's not lasting 4 hours at a time
- I find it hard to walk or talk
- I find it hard to breathe
- I'm coughing or wheezing a lot
- my chest is tight or it hurts.

If I have an asthma attack, I will:

1. Call for help. Sit up and try to keep calm.
2. Take 1 puff of my reliever inhaler (with my spacer, if I have it) every 30 to 60 seconds, up to a total of 10 puffs.
3. If I don't have my reliever inhaler, or it's not helping, or if I'm worried at any time, **call 999 for an ambulance.**
4. If the ambulance has not arrived after 10 minutes and my symptoms are not improving, repeat step 2.
5. If my symptoms are no better after repeating step 2, and the ambulance has still not arrived, **contact 999 again immediately.**

IMPORTANT: If you have a MART or AIR inhaler, please tell the responder when you **call 999.**

What to do after your child's asthma attack:

- If the asthma attack was managed at home, contact your GP surgery or call **111.**
- If your child was treated in hospital, take them to see a healthcare professional within 48 hours of being discharged.
- Make sure your child finishes any medicine they're prescribed, even if they start to feel better.
- If your child does not improve after treatment, see a healthcare professional urgently.